

Strong randomness of off-diagonal gluon phases and off-diagonal gluon mass in the maximally abelian gauge in QCD*

H. Suganuma^a, K. Amemiya^b, H. Ichie^c, N. Ishii^d, H. Matsufuru^e and T.T. Takahashi^f

^a Faculty of Science, Tokyo Institute of Technology, Ohokayama 2-12-1, Meguro, Tokyo 152-8551, Japan

^b Advanced Algorithm and Systems, Ebisu 1-13-6, Shibuya, Tokyo 150-0013, Japan

^c Humboldt Univ. zu Berlin, Institut für Physik, Invalidenstrasse 110, D-10115 Berlin, Germany

^d The Institute of Physical and Chemical Research (RIKEN), Hirosawa 2-1, Wako 351-0198, Japan

^e Yukawa Institute for Theoretical Physics (YITP), Kyoto University, Sakyo, Kyoto 606-8502, Japan

^f Research Center for Nuclear Physics, Osaka University, Mihogaoka 10-1, Ibaraki 567-0047, Japan

We study abelianization of QCD in the maximally abelian (MA) gauge. In the MA gauge, the off-diagonal gluon amplitude is strongly suppressed, and then the off-diagonal gluon phase shows strong randomness, which leads to a large off-diagonal gluon mass. Using lattice QCD, we find a large effective off-diagonal gluon mass in the MA gauge: $M_{\text{off}} \simeq 1.2\text{GeV}$ in SU(2) QCD, $M_{\text{off}} \simeq 1.1\text{GeV}$ in SU(3) QCD. Due to the large off-diagonal gluon mass in the MA gauge, infrared QCD is well abelianized like nonabelian Higgs theories. We investigate the inter-monopole potential and the dual gluon field B_μ in the MA gauge, and find longitudinal magnetic screening with $m_B \simeq 0.5\text{ GeV}$ in the infrared region, which indicates the dual Higgs mechanism by monopole condensation. We propose a gauge invariant description of the MA projection by introducing the “gluonic Higgs scalar field”.

1. Strong Randomness of Off-diagonal Gluon Phase and Large Mass of Off-diagonal Gluons in MA Gauge

In Euclidean QCD, the maximally abelian (MA) gauge is defined so as to minimize the total amount of the off-diagonal gluons [1–4]

$$R_{\text{off}}[A_\mu(\cdot)] \equiv \int d^4x \text{tr} \left\{ [\hat{D}_\mu, \vec{H}] [\hat{D}_\mu, \vec{H}]^\dagger \right\} \quad (1)$$

by the $SU(N_c)$ gauge transformation, from which the local MA gauge condition is easily derived as $[\vec{H}, [\hat{D}_\mu, [\hat{D}_\mu, \vec{H}]]] = 0$.

In SU(2) lattice QCD, we find two remarkable features of the off-diagonal gluon $A_\mu^\pm \equiv \frac{1}{\sqrt{2}}(A_\mu^1 \pm iA_\mu^2) = e^{\pm i\chi_\mu(x)} |A_\mu^\pm(x)|$ in the MA gauge [1–3].

1. The off-diagonal gluon amplitude $|A_\mu^\pm(x)|$ is strongly suppressed by $SU(N_c)$ gauge transformation in the MA gauge.
2. The off-diagonal gluon phase $\chi_\mu(x)$ tends to be random, because $\chi_\mu(x)$ is not con-

strained by MA gauge fixing at all, and only the constraint from the QCD action is weak due to a small accompanying factor $|A_\mu^\pm|$.

We investigate $\Delta\chi \equiv |\chi_\mu(s) - \chi_\mu(s + \hat{\nu})|(\text{mod } \pi)$ in the MA gauge with $U(1)_3$ Landau gauge fixing. If the off-diagonal gluon phase $\chi_\mu(x)$ is a continuum variable, as the lattice spacing a goes to 0, $\Delta\chi \simeq a|\partial_\nu \chi_\mu|$ goes to zero, and the probability distribution $P(\Delta\chi)$ approaches to $\delta(\Delta\chi)$. However, $P(\Delta\chi)$ is almost flat independently of a or β , which indicates *strong randomness of the off-diagonal gluon phase* $\chi_\mu(x)$ in the MA gauge.

Remarkably, *strong randomness of off-diagonal gluon phases* in the MA gauge leads to *rapid reduction of off-diagonal gluon correlations* [1] as

$$\begin{aligned} \langle A_\mu^+(x) A_\nu^-(y) \rangle &= \langle |A_\mu^+(x) A_\nu^-(y)| e^{i\{\chi_\mu(x) - \chi_\nu(y)\}} \rangle \\ &\sim \langle |A_\mu^\pm(x)|^2 \rangle_{\text{MA}} \delta_{\mu\nu} \delta^4(x - y), \quad (2) \end{aligned}$$

which means the infinitely large mass of off-diagonal gluons. Since the real off-diagonal gluon phases are not complete but approximate random phases even in the MA gauge, the off-diagonal gluon mass M_{off} would be large but finite.

*Talk presented by H. Suganuma

2. Large Mass Generation of Off-diagonal Gluons in MA Gauge : Essence of Infrared Abelianization of QCD

We study the Euclidean gluon propagator $G_{\mu\nu}^{ab}(x-y) \equiv \langle A_\mu^a(x) A_\nu^b(y) \rangle$ ($a, b = 1, 2, \dots, N_c^2 - 1$) and the off-diagonal gluon mass M_{off} in the MA gauge [1,3–5] using SU(N_c) lattice QCD with $N_c = 2, 3$. As for the residual abelian gauge symmetry, we take U(1)₃($\times$ U(1)₈) Landau gauge, to extract most continuous gluon configuration under the MA gauge constraint, for the comparison with the continuum theory. The continuum gluon field $A_\mu^a(x)$ is derived from the link variable as $U_\mu(s) = \exp\{iaeA_\mu^a(s)T^a\}$.

We show the scalar-type gluon propagators $G_{\mu\mu}^a(r) \equiv \langle A_\mu^a(x) A_\mu^a(y) \rangle$ ($a = 1, 2, \dots, N_c^2 - 1$), which depend only on the four-dimensional Euclidean distance $r \equiv \sqrt{(x_\mu - y_\mu)^2}$. The four-dimensional Euclidean propagator of the massive vector boson with the mass M takes a Yukawa-type asymptotic form as

$$G_{\mu\mu}(r) \simeq \frac{3}{4\pi^2} \frac{M}{r} K_1(Mr) \simeq \frac{3M^{1/2}}{2(2\pi)^{3/2}} \frac{e^{-Mr}}{r^{3/2}}. \quad (3)$$

From the slope analysis of the lattice QCD data of $\ln\{r^{3/2}G_{\mu\mu}(r)\}$ with $r \geq 0.2\text{fm}$, we obtain the off-diagonal gluon mass in the MA gauge as follows.

1. $M_{\text{off}} \simeq 1.2 \text{ GeV}$ in SU(2) lattice QCD² with $2.2 \leq \beta \leq 2.4$ and $12^3 \times 24, 16^4, 20^4$ (Fig.1).
2. $M_{\text{off}} \simeq 1.1 \text{ GeV}$ in SU(3) lattice QCD with $\beta = 5.7$ ($a \simeq 0.19\text{fm}$) and $12^3 \times 24$ (Fig.2).

In SU(3) QCD, the two diagonal gluon propagators, $G_{\mu\mu}^3$ and $G_{\mu\mu}^8$, show the same large distance propagation, and the three off-diagonal gluon propagators $G_{\mu\mu}^{+-(i,j)} \equiv \langle A_\mu^{+(i,j)}(x) A_\mu^{-(i,j)}(y) \rangle$ with $A_\mu^{\pm(i,j)} \equiv \frac{1}{\sqrt{2}}(A_\mu^i \pm iA_\mu^j)$ with $(i, j) = (1, 2), (4, 5), (6, 7)$ show the same massive behavior.

To conclude, both in SU(2) and SU(3) QCD, the off-diagonal gluon acquires a large effective mass $M_{\text{off}} \simeq 1\text{GeV}$ in the MA gauge, which is essence of infrared abelian dominance [1,3–8].

²From the mass measurement with the zero-momentum projection in SU(2) lattice QCD ($2.3 \leq \beta \leq 2.35$, $16^3 \times 32$, $12^3 \times 24$) we find again $M_{\text{off}} \simeq 1.2\text{GeV}$ in the MA gauge.




Figure 1. (a) The scalar-type gluon propagator $G_{\mu\mu}^a(r)$ v.s. 4-dim. distance r in the MA gauge in SU(2) lattice QCD with $2.2 \leq \beta \leq 2.4$, $12^3 \times 24$, 16^4 , 20^4 . (b) The logarithmic plot of $r^{3/2}G_{\mu\mu}^a(r)$.

3. Inter Monopole Potential, Longitudinal Magnetic Screening, Infrared Monopole Condensation and Monopole Structure

Using SU(2) lattice QCD, we study the *inter-monopole potential* and the *dual gluon propagator* in the monopole part in the MA gauge, and show *longitudinal magnetic screening* in the infrared region, as a direct evidence of the *dual Higgs mechanism by monopole condensation* [9,10]. The dual gluon mass is estimated as $m_B \simeq 0.5 \text{ GeV}$ [1,3,4]. Then, lattice QCD in the MA gauge exhibits *infrared abelian dominance* and *infrared monopole condensation*, which lead to the dual Ginzburg-Landau (DGL) theory [11] for infrared QCD.

Using SU(2) lattice QCD in the MA gauge, we find also the *monopole structure relating to a large amount of off-diagonal gluons around its center* like the 't Hooft-Polyakov monopole. At a large scale, off-diagonal gluons inside monopoles become invisible, and monopoles can be regarded as point-like Dirac monopoles [1–4].




Figure 2. (a) The scalar-type gluon propagator $G_{\mu\mu}^a(r)$ v.s. the four-dimensional distance r in the MA gauge with $U(1)_3 \times U(1)_8$ Landau gauge fixing in SU(3) lattice QCD with $\beta = 5.7$ and $12^3 \times 24$. (b) The logarithmic plot of $r^{3/2} G_{\mu\mu}^a(r)$.

4. Gluonic Higgs and Gauge Invariant Description of MA Projection

We propose a *gauge invariant description of the MA projection in QCD* [1,2] by introducing a “gluonic Higgs scalar field” $\vec{\phi}(x) \equiv \Omega(x) \vec{H} \Omega^\dagger(x)$ with $\Omega(x) \in \text{SU}(N_c)$ so as to minimize

$$R[\vec{\phi}(\cdot)] \equiv \int d^4x \text{tr} \left\{ [\hat{D}_\mu, \vec{\phi}(x)] [\hat{D}_\mu, \vec{\phi}(x)]^\dagger \right\} \quad (4)$$

for an arbitrary given gluon field $\{A_\mu(x)\}$.

The gluonic Higgs scalar $\vec{\phi}(x)$ physically corresponds to a “color-direction” of the nonabelian gauge connection $\hat{D}_\mu$ averaged over μ at each x .

Through the projection along $\vec{\phi}(x)$, we can extract the abelian $U(1)^{N_c-1}$ sub-gauge-manifold which is most close to the original $\text{SU}(N_c)$ gauge manifold. This projection is manifestly gauge invariant, and is mathematically equivalent to the ordinary MA projection [1,2]. In this description, we find *infrared relevance of the gluon mode along*

the color-direction $\vec{\phi}(x)$ [1,2,12], corresponding to infrared abelian dominance in the MA gauge.

Similar to $\hat{D}_\mu$, the gluonic Higgs scalar $\vec{\phi}(x)$ obeys the adjoint gauge transformation, and $\vec{\phi}(x)$ is diagonalized in the MA gauge. Then, *monopoles appear at the hedgehog singularities of $\vec{\phi}(x)$* , when the gluon field is continuous as in the SU(2) Landau gauge as shown in Fig.3 [1,2].




Figure 3. The correlation between the gluonic Higgs scalar field $\phi(x) = \phi^a(x) \frac{\tau^a}{2}$ and monopoles (dots) in the SU(2) Landau gauge in SU(2) lattice QCD with $\beta = 2.4$ and 16^4 . The arrow denotes the SU(2) color direction of $(\phi^1(x), \phi^2(x), \phi^3(x))$. The monopole appears at the hedgehog singularity of the gluonic Higgs scalar $\phi(x)$.

REFERENCES

1. H. Suganuma et al., *Quantum Chromodynamics and Color Confinement*, edited by H. Suganuma et al. (World Scientific, 2001) p.103.
2. H. Ichie, H. Suganuma, Phys.Rev.D60 (1999) 77501; Nucl.Phys.B548 (99) 365; B574 (00) 70.
3. H. Suganuma, K. Amemiya, H. Ichie and A. Tanaka, Nucl. Phys. A670 (2000) 40.
4. H.Suganuma, H.Ichie, A.Tanaka, K.Amemiya Prog. Theor. Phys. Suppl. 131 (1998) 559.
5. K. Amemiya and H. Suganuma, Phys. Rev. D60 (1999) 114509.
6. Z.F. Ezawa and A. Iwazaki, Phys. Rev. D25 (1982) 2681; D26 (1982) 631.
7. O. Miyamura, Phys. Lett. B353 (1995) 91.
8. K.-I. Kondo, Phys.Rev.D57 (1998) 7467; D58 (1998) 105016; Phys. Lett. B455 (1999) 251.
9. Y. Nambu, Phys. Rev. D10 (1974) 4262.
10. G. 't Hooft, Nucl. Phys. B190 (1981) 455.
11. H. Suganuma, S. Sasaki and H. Toki, Nucl. Phys. B435 (1995) 207.
12. Y.M. Cho, Phys. Rev. D21 (1980) 1080; D62 (2000) 074009.